

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 8 \cdot \frac{1}{1+3} \cdot x^{1+3} + C$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 8 \cdot \frac{1}{4} \cdot x^4 + C$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 2 \cdot x^4 + C$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 2 \cdot x^4 + C$$

$$\int (-e^x) dx =$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 2 \cdot x^4 + C$$

$$\int (-e^x) dx = -e^x + C$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 2 \cdot x^4 + C$$

$$\int (-e^x) dx = -e^x + C$$

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 2 \cdot x^4 + C$$

$$\int (-e^x) dx = -e^x + C$$

$$\int_0^1 (8 \cdot x^3 - e^x) dx = [2 \cdot x^4 - e^x]_0^1 =$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 2 \cdot x^4 + C$$

$$\int (-e^x) dx = -e^x + C$$

$$\begin{aligned} \int_0^1 (8 \cdot x^3 - e^x) dx &= \left[2 \cdot x^4 - e^x \right]_0^1 = \\ (2 \cdot 1^4 - e^1) - (2 \cdot 0^4 - e^0) &= \end{aligned}$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 2 \cdot x^4 + C$$

$$\int (-e^x) dx = -e^x + C$$

$$\begin{aligned} \int_0^1 (8 \cdot x^3 - e^x) dx &= [2 \cdot x^4 - e^x]_0^1 = \\ (2 \cdot 1^4 - e^1) - (2 \cdot 0^4 - e^0) &= (2 - e) - (-1) = \end{aligned}$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)

Bestem integralet

$$\int_0^1 (8 \cdot x^3 - e^x) dx$$

Hvert led integreres for sig.

$$\int (8 \cdot x^3) dx = 2 \cdot x^4 + C$$

$$\int (-e^x) dx = -e^x + C$$

$$\begin{aligned} \int_0^1 (8 \cdot x^3 - e^x) dx &= [2 \cdot x^4 - e^x]_0^1 = \\ (2 \cdot 1^4 - e^1) - (2 \cdot 0^4 - e^0) &= (2 - e) - (-1) = 3 - e \end{aligned}$$

$f(x)$	$F(x)$	
k	$k \cdot x + C$	(1)
$k \cdot x$	$k \cdot \frac{1}{2} \cdot x^2 + C$	(2)
x^n	$\frac{1}{n+1} \cdot x^{n+1} + C$	(3)
e^x	$e^x + C$	(4)
$\frac{1}{x}$	$\ln(x) + C$	(5)
\sqrt{x}	$\frac{2}{3}x^{3/2} + C$	(6)