

Differentialkvotient for $f(x) = g(h(x))$

**Differentialkvotient for $f(x) = g(h(x))$,
hvor g og h er en differentiable funktioner.**

Trin 1: Indsæt f i differenskvotienten.

Differentialkvotient for $f(x) = g(h(x))$

**Differentialkvotient for $f(x) = g(h(x))$,
hvor g og h er en differentiable funktioner.**

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$,
hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$\frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = v$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$,
hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$\frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} = v + g'(y_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$,
hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$g(y_0 + \mathbf{k}) - g(y_0) = (v + g'(y_0)) \cdot \mathbf{k}$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$,
hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$,
hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$
$$h(x_0 + \mathbf{h}) = (w + h'(x_0)) \cdot \mathbf{h} + h(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$

$$h(x_0 + \mathbf{h}) = (w + h'(x_0)) \cdot \mathbf{h} + h(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$\frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$

$$h(x_0 + \mathbf{h}) = (w + h'(x_0)) \cdot \mathbf{h} + h(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$\frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}} = \frac{g((w + h'(x_0)) \cdot \mathbf{h} + h(x_0)) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$
$$h(x_0 + \mathbf{h}) = (w + h'(x_0)) \cdot \mathbf{h} + h(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$
$$h(x_0 + \mathbf{h}) = (w + h'(x_0)) \cdot \mathbf{h} + h(x_0)$$
$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

Trin 2: Reducer differenskvotienten.

$$\frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}} = \frac{g(h(x_0) + (w + h'(x_0)) \cdot \mathbf{h}) - g(h(x_0))}{\mathbf{h}}$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$\frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}} = \frac{(v + g'(h(x_0))) \cdot (w + h'(x_0)) \cdot \mathbf{h} + g(h(x_0)) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$

$$h(x_0 + \mathbf{h}) = (w + h'(x_0)) \cdot \mathbf{h} + h(x_0)$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$\frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}} = \frac{(v + g'(h(x_0))) \cdot (w + h'(x_0)) \cdot \mathbf{h}}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$

$$h(x_0 + \mathbf{h}) = (w + h'(x_0)) \cdot \mathbf{h} + h(x_0)$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$\frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}} = (v + g'(h(x_0))) \cdot (w + h'(x_0))$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$g(y_0 + \mathbf{k}) = (v + g'(y_0)) \cdot \mathbf{k} + g(y_0)$$

$$h(x_0 + \mathbf{h}) = (w + h'(x_0)) \cdot \mathbf{h} + h(x_0)$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$
$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$\lim_{\mathbf{h} \rightarrow 0} (v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$\lim_{\mathbf{h} \rightarrow 0} (v + g'(h(x_0))) \cdot (w + h'(x_0))$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

$$v = \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$\lim_{\mathbf{h} \rightarrow 0} (v + g'(h(x_0))) \cdot (w + h'(x_0))$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

$$v = \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0)$$

$$\lim_{\mathbf{h} \rightarrow 0} \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0) = 0$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$\lim_{\mathbf{h} \rightarrow 0} (v + g'(h(x_0))) \cdot (w + h'(x_0))$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

$$v = \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0)$$

$$\lim_{\mathbf{h} \rightarrow 0} \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0) = 0$$

$$w = \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$\left(\lim_{\mathbf{h} \rightarrow 0} v + \lim_{\mathbf{h} \rightarrow 0} g'(h(x_0)) \right) \cdot \left(\lim_{\mathbf{h} \rightarrow 0} w + \lim_{\mathbf{h} \rightarrow 0} h'(x_0) \right)$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

$$v = \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0)$$

$$\lim_{\mathbf{h} \rightarrow 0} \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0) = 0$$

$$w = \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$\left(\lim_{\mathbf{h} \rightarrow 0} v + g'(h(x_0)) \right) \cdot \left(\lim_{\mathbf{h} \rightarrow 0} w + h'(x_0) \right)$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

$$v = \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0)$$

$$\lim_{\mathbf{h} \rightarrow 0} \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0) = 0$$

$$w = \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$\left(\lim_{\mathbf{h} \rightarrow 0} v + g'(h(x_0)) \right) \cdot h'(x_0)$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

$$v = \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0)$$

$$\lim_{\mathbf{h} \rightarrow 0} \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0) = 0$$

$$w = \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$g'(h(x_0)) \cdot h'(x_0)$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

$$v = \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0)$$

$$\lim_{\mathbf{h} \rightarrow 0} \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0) = 0$$

$$w = \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0)$$

Differentialkvotient for $f(x) = g(h(x))$

Differentialkvotient for $f(x) = g(h(x))$, hvor g og h er en differentiable funktioner.

Trin 1: Indsæt f i differenskvotienten.

$$\frac{f(x_0 + \mathbf{h}) - f(x_0)}{\mathbf{h}} = \frac{g(h(x_0 + \mathbf{h})) - g(h(x_0))}{\mathbf{h}}$$

Trin 2: Reducer differenskvotienten.

$$(v + g'(h(x_0))) \cdot (w + h'(x_0))$$

Trin 3: Udregn grænseværdien.

$$f'(x) = g'(h(x)) \cdot h'(x)$$

$$\lim_{\mathbf{k} \rightarrow 0} \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0) = 0$$

$$\mathbf{k} = (w + h'(x_0)) \cdot \mathbf{h}$$

$$v = \frac{g(y_0 + \mathbf{k}) - g(y_0)}{\mathbf{k}} - g'(y_0)$$

$$\lim_{\mathbf{h} \rightarrow 0} \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0) = 0$$

$$w = \frac{h(x_0 + \mathbf{h}) - h(x_0)}{\mathbf{h}} - h'(x_0)$$

Anvendelse af regneregler

Bestem f' for følgende funktioner.

$$f(x) = \sqrt{2x^2 + 4x}$$

$$f(x) = \ln(x^2 + 3)$$

$$f(x) = e^{x^2+1}$$

f	f'	
$g + h$	$g' + h'$	(10)
$k \cdot g(x)$	$k \cdot g'(x)$	(11)
$g \cdot h$	$g' \cdot h + g \cdot h'$	(12)
$\frac{1}{g}$	$-\frac{g'}{g^2}$	(13)
$\frac{h}{g}$	$\frac{h' \cdot g - h \cdot g'}{g^2}$	(14)
$g(h(x))$	$g'(h(x)) \cdot h'(x)$	(15)

f	f'	
k	0	(1)
$k \cdot x$	k	(2)
x^n	$n \cdot x^{n-1}$	(3)
$\frac{1}{x}$	$-\frac{1}{x^2}$	(4)
\sqrt{x}	$\frac{1}{2\sqrt{x}}$	(5)
e^x	e^x	(6)
$e^{k \cdot x}$	$k \cdot e^{k \cdot x}$	(7)
$\ln(x)$	$\frac{1}{x}$	(8)
a^x	$a^x \cdot \ln(a)$	(9)

Anvendelse af regneregler

Bestem f' for følgende funktioner.

$$f(x) = \sqrt{2x^2 + 4x} \Rightarrow f'(x) = \frac{1}{2\sqrt{2x^2 + 4x}} \cdot (4x + 4)$$

$$f(x) = \ln(x^2 + 3)$$

$$f(x) = e^{x^2+1}$$

f	f'	
$g + h$	$g' + h'$	(10)
$k \cdot g(x)$	$k \cdot g'(x)$	(11)
$g \cdot h$	$g' \cdot h + g \cdot h'$	(12)
$\frac{1}{g}$	$-\frac{g'}{g^2}$	(13)
$\frac{h}{g}$	$\frac{h' \cdot g - h \cdot g'}{g^2}$	(14)
$g(h(x))$	$g'(h(x)) \cdot h'(x)$	(15)

f	f'	
k	0	(1)
$k \cdot x$	k	(2)
x^n	$n \cdot x^{n-1}$	(3)
$\frac{1}{x}$	$-\frac{1}{x^2}$	(4)
\sqrt{x}	$\frac{1}{2\sqrt{x}}$	(5)
e^x	e^x	(6)
$e^{k \cdot x}$	$k \cdot e^{k \cdot x}$	(7)
$\ln(x)$	$\frac{1}{x}$	(8)
a^x	$a^x \cdot \ln(a)$	(9)

Anvendelse af regneregler

Bestem f' for følgende funktioner.

$$f(x) = \sqrt{2x^2 + 4x} \Rightarrow f'(x) = \frac{(4x + 4)}{2\sqrt{2x^2 + 4x}}$$

$$f(x) = \ln(x^2 + 3)$$

$$f(x) = e^{x^2+1}$$

f	f'	
$g + h$	$g' + h'$	(10)
$k \cdot g(x)$	$k \cdot g'(x)$	(11)
$g \cdot h$	$g' \cdot h + g \cdot h'$	(12)
$\frac{1}{g}$	$-\frac{g'}{g^2}$	(13)
$\frac{h}{g}$	$\frac{h' \cdot g - h \cdot g'}{g^2}$	(14)
$g(h(x))$	$g'(h(x)) \cdot h'(x)$	(15)

f	f'	
k	0	(1)
$k \cdot x$	k	(2)
x^n	$n \cdot x^{n-1}$	(3)
$\frac{1}{x}$	$-\frac{1}{x^2}$	(4)
\sqrt{x}	$\frac{1}{2\sqrt{x}}$	(5)
e^x	e^x	(6)
$e^{k \cdot x}$	$k \cdot e^{k \cdot x}$	(7)
$\ln(x)$	$\frac{1}{x}$	(8)
a^x	$a^x \cdot \ln(a)$	(9)

Anvendelse af regneregler

Bestem f' for følgende funktioner.

$$f(x) = \sqrt{2x^2 + 4x} \Rightarrow f'(x) = \frac{2(2x + 2)}{2\sqrt{2x^2 + 4x}}$$

$$f(x) = \ln(x^2 + 3)$$

$$f(x) = e^{x^2+1}$$

f	f'	
$g + h$	$g' + h'$	(10)
$k \cdot g(x)$	$k \cdot g'(x)$	(11)
$g \cdot h$	$g' \cdot h + g \cdot h'$	(12)
$\frac{1}{g}$	$-\frac{g'}{g^2}$	(13)
$\frac{h}{g}$	$\frac{h' \cdot g - h \cdot g'}{g^2}$	(14)
$g(h(x))$	$g'(h(x)) \cdot h'(x)$	(15)

f	f'	
k	0	(1)
$k \cdot x$	k	(2)
x^n	$n \cdot x^{n-1}$	(3)
$\frac{1}{x}$	$-\frac{1}{x^2}$	(4)
\sqrt{x}	$\frac{1}{2\sqrt{x}}$	(5)
e^x	e^x	(6)
$e^{k \cdot x}$	$k \cdot e^{k \cdot x}$	(7)
$\ln(x)$	$\frac{1}{x}$	(8)
a^x	$a^x \cdot \ln(a)$	(9)

Anvendelse af regneregler

Bestem f' for følgende funktioner.

$$f(x) = \sqrt{2x^2 + 4x} \Rightarrow f'(x) = \frac{(2x + 2)}{\sqrt{2x^2 + 4x}}$$

$$f(x) = \ln(x^2 + 3)$$

$$f(x) = e^{x^2+1}$$

f	f'	
$g + h$	$g' + h'$	(10)
$k \cdot g(x)$	$k \cdot g'(x)$	(11)
$g \cdot h$	$g' \cdot h + g \cdot h'$	(12)
$\frac{1}{g}$	$-\frac{g'}{g^2}$	(13)
$\frac{h}{g}$	$\frac{h' \cdot g - h \cdot g'}{g^2}$	(14)
$g(h(x))$	$g'(h(x)) \cdot h'(x)$	(15)

f	f'	
k	0	(1)
$k \cdot x$	k	(2)
x^n	$n \cdot x^{n-1}$	(3)
$\frac{1}{x}$	$-\frac{1}{x^2}$	(4)
\sqrt{x}	$\frac{1}{2\sqrt{x}}$	(5)
e^x	e^x	(6)
$e^{k \cdot x}$	$k \cdot e^{k \cdot x}$	(7)
$\ln(x)$	$\frac{1}{x}$	(8)
a^x	$a^x \cdot \ln(a)$	(9)

Anvendelse af regneregler

Bestem f' for følgende funktioner.

$$f(x) = \sqrt{2x^2 + 4x} \Rightarrow f'(x) = \frac{(2x + 2)}{\sqrt{2x^2 + 4x}}$$

$$f(x) = \ln(x^2 + 3) \Rightarrow f'(x) = \frac{1}{x^2 + 3} \cdot 2x$$

$$f(x) = e^{x^2+1}$$

f	f'	
$g + h$	$g' + h'$	(10)
$k \cdot g(x)$	$k \cdot g'(x)$	(11)
$g \cdot h$	$g' \cdot h + g \cdot h'$	(12)
$\frac{1}{g}$	$-\frac{g'}{g^2}$	(13)
$\frac{h}{g}$	$\frac{h' \cdot g - h \cdot g'}{g^2}$	(14)
$g(h(x))$	$g'(h(x)) \cdot h'(x)$	(15)

f	f'	
k	0	(1)
$k \cdot x$	k	(2)
x^n	$n \cdot x^{n-1}$	(3)
$\frac{1}{x}$	$-\frac{1}{x^2}$	(4)
\sqrt{x}	$\frac{1}{2\sqrt{x}}$	(5)
e^x	e^x	(6)
$e^{k \cdot x}$	$k \cdot e^{k \cdot x}$	(7)
$\ln(x)$	$\frac{1}{x}$	(8)
a^x	$a^x \cdot \ln(a)$	(9)

Anvendelse af regneregler

Bestem f' for følgende funktioner.

$$f(x) = \sqrt{2x^2 + 4x} \Rightarrow f'(x) = \frac{(2x + 2)}{\sqrt{2x^2 + 4x}}$$

$$f(x) = \ln(x^2 + 3) \Rightarrow f'(x) = \frac{2x}{x^2 + 3}$$

$$f(x) = e^{x^2+1}$$

f	f'	
$g + h$	$g' + h'$	(10)
$k \cdot g(x)$	$k \cdot g'(x)$	(11)
$g \cdot h$	$g' \cdot h + g \cdot h'$	(12)
$\frac{1}{g}$	$-\frac{g'}{g^2}$	(13)
$\frac{h}{g}$	$\frac{h' \cdot g - h \cdot g'}{g^2}$	(14)
$g(h(x))$	$g'(h(x)) \cdot h'(x)$	(15)

f	f'	
k	0	(1)
$k \cdot x$	k	(2)
x^n	$n \cdot x^{n-1}$	(3)
$\frac{1}{x}$	$-\frac{1}{x^2}$	(4)
\sqrt{x}	$\frac{1}{2\sqrt{x}}$	(5)
e^x	e^x	(6)
$e^{k \cdot x}$	$k \cdot e^{k \cdot x}$	(7)
$\ln(x)$	$\frac{1}{x}$	(8)
a^x	$a^x \cdot \ln(a)$	(9)

Anvendelse af regneregler

Bestem f' for følgende funktioner.

$$f(x) = \sqrt{2x^2 + 4x} \Rightarrow f'(x) = \frac{(2x + 2)}{\sqrt{2x^2 + 4x}}$$

$$f(x) = \ln(x^2 + 3) \Rightarrow f'(x) = \frac{2x}{x^2 + 3}$$

$$f(x) = e^{x^2+1} \Rightarrow f'(x) = e^{x^2+1} \cdot 2x$$

f	f'	
$g + h$	$g' + h'$	(10)
$k \cdot g(x)$	$k \cdot g'(x)$	(11)
$g \cdot h$	$g' \cdot h + g \cdot h'$	(12)
$\frac{1}{g}$	$-\frac{g'}{g^2}$	(13)
$\frac{h}{g}$	$\frac{h' \cdot g - h \cdot g'}{g^2}$	(14)
$g(h(x))$	$g'(h(x)) \cdot h'(x)$	(15)

f	f'	
k	0	(1)
$k \cdot x$	k	(2)
x^n	$n \cdot x^{n-1}$	(3)
$\frac{1}{x}$	$-\frac{1}{x^2}$	(4)
\sqrt{x}	$\frac{1}{2\sqrt{x}}$	(5)
e^x	e^x	(6)
$e^{k \cdot x}$	$k \cdot e^{k \cdot x}$	(7)
$\ln(x)$	$\frac{1}{x}$	(8)
a^x	$a^x \cdot \ln(a)$	(9)