

Symmetri af differentiationsrækkefølge

12. april 2020

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a)$$

$$Q = v(b + k) - v(b)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a)$$

$$Q = v(b + k) - v(b)$$

$$\frac{u(a + h) - u(a)}{(a + h) - a} = u'(a + i_1 \cdot h), \quad 0 \leq i_1 < 1$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a)$$

$$Q = v(b + k) - v(b)$$

$$\frac{u(a + h) - u(a)}{(a + h) - a} = u'(a + i_1 \cdot h), \quad 0 \leq i_1 < 1$$

$$\frac{u(a + h) - u(a)}{h} = u'(a + i_1 \cdot h)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a)$$

$$Q = v(b + k) - v(b)$$

$$\frac{u(a + h) - u(a)}{(a + h) - a} = u'(a + i_1 \cdot h), \quad 0 \leq i_1 < 1$$

$$\frac{u(a + h) - u(a)}{h} = u'(a + i_1 \cdot h)$$

$$u(a + h) - u(a) = h \cdot u'(a + i_1 \cdot h)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h)$$

$$Q = v(b + k) - v(b)$$

$$\frac{u(a + h) - u(a)}{(a + h) - a} = u'(a + i_1 \cdot h), \quad 0 \leq i_1 < 1$$

$$\frac{u(a + h) - u(a)}{h} = u'(a + i_1 \cdot h)$$

$$u(a + h) - u(a) = h \cdot u'(a + i_1 \cdot h)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h) \\ = h \cdot (f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b))$$

$$Q = v(b + k) - v(b)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h) \\ = h \cdot (f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b))$$

$$Q = v(b + k) - v(b)$$

$$\frac{f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)}{(b + k) - b} \\ = f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h) \\ = h \cdot (f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b))$$

$$Q = v(b + k) - v(b)$$

$$\frac{f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)}{(b + k) - b} \\ = f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)$$

$$\frac{f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)}{k} \\ = f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h) \\ = h \cdot (f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b))$$

$$Q = v(b + k) - v(b)$$

$$\frac{f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)}{(b + k) - b} \\ = f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)$$

$$\frac{f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)}{k} \\ = f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)$$

$$f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b) \\ = k \cdot f''_{xy}(a + i_1 \cdot h, b + i \cdot k)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h) \\ = h \cdot (f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)) \\ = h \cdot (k \cdot f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)) \\ Q = v(b + k) - v(b)$$

$$\frac{f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)}{(b + k) - b} \\ = f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)$$

$$\frac{f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)}{k} \\ = f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)$$

$$f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b) \\ = k \cdot f''_{xy}(a + i_1 \cdot h, b + i \cdot k)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h) \\ = h \cdot (f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)) \\ = h \cdot (k \cdot f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k))$$

$$Q = v(b + k) - v(b) \\ = k \cdot (h \cdot f''_{yx}(a + i_4 \cdot h, b + i_3 \cdot k))$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h) \\ = h \cdot (f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)) \\ = h \cdot (k \cdot f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)) \\ Q = v(b + k) - v(b) \\ = k \cdot (h \cdot f''_{yx}(a + i_4 \cdot h, b + i_3 \cdot k))$$

$$f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k) = f''_{yx}(a + i_3 \cdot h, b + i_4 \cdot k)$$

Bevis for at $f''_{xy}(x, y) = f''_{yx}(x, y)$
i $P(a, b, f(a, b))$

$$Q = f(a + h, b + k) - f(a + h, b) \\ - f(a, b + k) + f(a, b)$$

$$u(x) = f(x, b + k) - f(x, b)$$

$$v(y) = f(a + h, y) - f(a, y)$$

$$Q = u(a + h) - u(a) \\ = h \cdot u'(a + i_1 \cdot h) \\ = h \cdot (f'_x(a + i_1 \cdot h, b + k) - f'_x(a + i_1 \cdot h, b)) \\ = h \cdot (k \cdot f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k)) \\ Q = v(b + k) - v(b) \\ = k \cdot (h \cdot f''_{yx}(a + i_4 \cdot h, b + i_3 \cdot k))$$

$$f''_{xy}(a + i_1 \cdot h, b + i_2 \cdot k) = f''_{yx}(a + i_3 \cdot h, b + i_4 \cdot k) \\ f''_{xy}(a, b) = f''_{yx}(a, b)$$